

Papéis Avulsos de Zoologia



This is an open-access article distributed under the terms of the Creative Commons Attribution License. Fonte:

<https://www.scielo.br/j/paz/a/CKgQNsMN43s8fYtGnWGpWsb/?lang=en#>. Acesso em: 10 nov. 2021.

REFERÊNCIA

HARTERREITEN-SOUZA, Érica Sevilha *et al.* Seasonal fluctuation in the population of *Harmonia axyridis* (Pallas, 1773) (Coleoptera: Coccinellidae) and co-occurrence with other Coccinellids in the Federal District of Brazil. **Papéis Avulsos de Zoologia**, v. 52, n. 11, p. 134-140, 2012. DOI: <https://doi.org/10.1590/S0031-10492012001100001>.

Disponível em:

<https://www.scielo.br/j/paz/a/CKgQNsMN43s8fYtGnWGpWsb/?lang=en#>. Acesso em: 10 nov. 2021.

Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 52(11):133-139, 2012

www.mz.usp.br/publicacoes
<http://portal.revistasusp.sibi.usp.br>
www.scielo.br/paz

ISSN impresso: 0031-1049

ISSN on-line: 1807-0205

SEASONAL FLUCTUATION IN THE POPULATION OF *HARMONIA AXYRIDIS* (PALLAS, 1773) (COLEOPTERA: COCCINELLIDAE) AND CO-OCCURRENCE WITH OTHER COCCINELLIDS IN THE FEDERAL DISTRICT OF BRAZIL

ÉRICA SEVILHA HARTERREITEN-SOUZA¹
PEDRO HENRIQUE BRUM TOGNI²
PALOMA VIRGÍNIA GAMBARRA NITÁO MILANE³
KELLY RAMALHO CAVALCANTE⁵
MARIA ALICE DE MEDEIROS⁴
CARMEN SILVIA SOARES PIRES⁵
EDISON RYOITI SUJII⁵

ABSTRACT

The multicolored Asian lady beetle, *Harmonia axyridis* (Pallas, 1773), was first recorded in Brazil in 2002 in Paraná state and subsequently observed in Santa Catarina, Rio Grande do Sul and Minas Gerais. This species can spread to new areas and become dominant in the local community, reducing the density and diversity of native species, mainly Coccinellidae. The objective of this work was to record for the first time the occurrence of *H. axyridis* in the Federal District and its co-occurrence with other Coccinellidae species. The individuals were collected directly from plants at an organic farm in Taguatinga and in experimental fields of Embrapa Hortaliças, located in the Federal District, from August 2008 to January 2010. We collected 881 Coccinellids, and of these, 110 belong to the species *H. axyridis*. These were found exclusively on the following plants of the succinea group: maize, cabbage, cauliflower, broccoli, kale, cucumber, cotton, tomato and coriander. We also observed its co-occurrence with the following lady beetle species: *Cycloneda sanguinea* (Linnaeus, 1763), *Hippodamia convergens* (Guerin-Meneville 1842), *Eriopis connexa* (German, 1824), *Scymnus* sp., *Nephaspis* sp., *Azya luteipes* (Mulsant, 1850), *Hyperaspis festiva* (Mulsant, 1850), *Olla v-nigrum* (Mulsant, 1866), *Psyllobora* sp. and *Coleomegilla maculata* (De Geer, 1775). So far, we have

1. Department of Ecology, Institute of Biological Sciences, Universidade de Brasília, Campus Darcy Ribeiro, 70910-900, Brasília, Distrito Federal, Brasil. E-mail: ericsevilha@hotmail.com
2. Department of Animal Biology/Entomology, Universidade Federal de Viçosa, Avenida PH Holfs, s/nº, Campus Universitário, 36570-000, Viçosa, MG, Brasil.
3. Department Agroecology and Environment. Empresa de Assistência Técnica e Extensão Rural do Distrito Federal, EMATER-DF. Parque Estação Biológica, Asa Norte, s/nº, 70770-915, Brasília, DF, Brasil.
4. Empresa Brasileira de Pesquisa Agropecuária, EMBRAPA Hortaliças. Rodovia BR-060 (Brasília-Anápolis), km 9, Fazenda Tamanduá, Ponte Alta, 70359-970, Gama, DF, Brasil.
5. Empresa Brasileira de Pesquisa Agropecuária, EMBRAPA Recursos Genéticos e Biotecnologia. Parque Estação Biológica, Asa Norte, 70770-917, Brasília, DF, Brasil.

not found any negative interactions between *H. axyridis* and these species. This is the northernmost *H. axyridis* record in Brazil. Moreover, the region was previously considered to have a low probability of occurrence for this species. Therefore, this record confirms that *H. axyridis* presents great adaptive plasticity to new habitats.

KEY-WORDS: Biological Control; Diversity; Exotic species; Invasion; Predator.

INTRODUCTION

The lady beetle *Harmonia axyridis* is an introduced species of Coccinellidae originally from the Palaearctic region (Koch, 2003). It is a voracious predator of aphids, also feeding on other insects such as psilids (Koch, 2003), eggs of Lepidoptera (Santos *et al.*, 2009) and pollen (Berkvens *et al.*, 2008). Due to this, *H. axyridis* has been used successfully in several biological control programs worldwide (Koch *et al.*, 2006). In Europe (Brown *et al.*, 2008), China (Zhang, 1992), the United States and Mexico (Brown & Miller, 1998), for example, the species is already widespread and is considered one of the main agents for biological control of aphids on different crops. The current widespread distribution of this species is related to its natural dispersal (Osawa, 2000), dispersal facilitated by human action (Brown *et al.*, 2008), and adaptability to climates different from its original region (Koch *et al.*, 2006).

However, the characteristics that make *H. axyridis* an efficient predator and a good biological control agent have also led to it becoming a threat to native guilds of aphid-eaters in regions where it has been introduced. *H. axyridis* is bigger than other coccinellids, has high predatory capacity (Elliott *et al.*, 1996), high fecundity (Iablokoff-Khnzorian, 1982), and can act as an intraguild predator mainly on other coccinellids (*e.g.*, Burgio *et al.*, 2003, Santos *et al.*, 2009). This means that it can displace several native aphid predators, becoming dominant in the local community, as observed in southern Brazil (Martins *et al.*, 2009).

H. axyridis was introduced in the early 1990s in South America through the province of Mendoza in Argentina as a means of biologically controlling aphids on peaches (Saini, 2004). Currently there are reports of this species in Chile (Grez *et al.*, 2010) and Perú (González, 2007). In Brazil, it was first recorded in Curitiba, Paraná state, in April 2002 feeding on *Tinocallis kahawaluokalani* (Kirkaldy) (Hemiptera: Aphididae), *Lagerstroemia indica* (Linnaeus) (Lythraceae), *Cinara* spp. (Hemiptera: Aphididae), and *Pinus* spp. (Pinaceae) (Almeida & Silva, 2002). In 2006 it was recorded in Viçosa, Minas Gerais, feeding on aphids and psyllids in ornamental plants and

vegetables (Rezende *et al.*, 2010). Although the presence of this species is recent in Brazil, significant reduction in the abundance of other native species of coccinellids has been recorded (Milléo *et al.*, 2008), demonstrating the potential threat of this species to the guild of aphid predators in Brazil (Martins *et al.*, 2009).

Due to the threat that this species may represent to predators of the local insect fauna, such as coccinellids, the objective of this study was to evaluate seasonal variation in the abundance of *H. axyridis* and other species of Coccinellidae on some organic vegetable crops in the Federal District. This would make possible the evaluation of shifts in richness and diversity of *H. axyridis* at an early point of establishment in the region.

MATERIAL AND METHODS

Sampling was done on plantations of organic vegetables in the Federal District from August 2008 to January 2010. In order to collect natural enemies of tomato, initial surveys were performed on tomatoes and cilantro crops in the experimental field of Embrapa Hortaliças Research Center in the region of Gama (15°56'S and 48°08'W). Samples were taken weekly from August to November 2008 (end of drought in the region) by directly collecting individuals of Coccinellidae from 240 randomly designed tomato plants. The first record of *H. axyridis* was in 2008 at Embrapa Hortaliças and it was collected again in 2009 in an intercropping area of vegetables (corn, cabbage, kale, cauliflower, cucumbers, broccoli, tomato and cilantro) at an organic farm in Taguatinga (15°49'S and 48°04'W). After that, monitoring of Coccinellidae species population at this property was done monthly from February 2009 to January 2010. Collection of insects was made directly from plants with a sampling effort of two hours.

Different patterns of color and spot distribution were found for *H. axyridis* occurring in the Federal District. Some specimens of each pattern were mounted, photographed and sent for identification by a specialist (Dr. Lúcia Massuti from the

Universidade Federal do Paraná). Specimens were deposited in the Entomological PE. J.S. Moure, Department of Zoology, Universidade Federal do Paraná, where they were numbered sequentially from DZUP/144661 DZUP/144676. The remaining specimens were compared and deposited in the Insect Collection at Embrapa Recursos Genéticos e Biotecnologia.

Abundance of *H. axyridis* and other species of lady beetle was recorded for each date, place and culture in order to assess the community structure of coccinellids by comparing the relative abundances of *H. axyridis* in relation to the total abundance of species. Species richness of each area was estimated by rarefaction curves. Diversity was compared by t test, calculating the Shannon-Wiener diversity index and

Renyi profile index through use of the statistical program PAST (Hammer *et al.*, 2001).

RESULTS

We collected 881 coccinellids, with 110 of these belonging to the species *H. axyridis* exclusively in the succinae group. Color ranged from red to orange with a variable number of black spots or complete lack thereof (Fig. 1). Other coccinellid species that co-occurred in the two sampling areas were *Azya luteipes*, *Coleomegilla maculate*, *Cycloneda sanguinea*, *Eriopis connexa*, *Hippodamia convergens*, *Hyperaspis festiva*, *Nephaspis* sp., *Olla v-nigrum*, *Psyllobora* spp. and *Scymnus* spp. (Table 1).




FIGURE 1: Coloration patterns of *Harmonia axyridis* collected in the region of the Federal District, Brazil. The scale bar above the photos of the specimens corresponds to 4 mm.


Only one specimen of *Harmonia axyridis* was collected in mixed stands of tomato and cilantro in Gama. The most abundant species collected from the local community of coccinellids were *Hippodamia convergens*, *Eriopis connexa*, *Scymnus* spp. and *Cycloneda sanguinea*. In the intercropping of vegetables in Taguatinga, however, *H. axyridis* was the dominant species in the local community of coccinellids (Table 1). On this property, the proportion of *H. axyridis* compared with other coccinellid species accounted

for 39.4% of specimens (Table 1). Although the climate of the region shows a strong variation in rainfall (Fig. 2), specimens of *H. axyridis* were collected both in the rainy season and during the dry season, with higher peaks of abundance during the dry season in August and October (Fig. 3).

The comparison of coccinellid communities collected from the two areas showed the same number of species (Table 1), although the richness estimated by rarefaction curves (Fig. 4) was higher in Taguatinga.



A Rainfall Higher temperature Lower temperature



B Rainfall Higher temperature Lower temperature

FIGURE 2: Seasonal pattern of rainfall distribution and variation in mean maximum and minimum temperatures in the Federal District, (A) 2008 and (B) 2009.

TABLE 1: Relative abundance of coccinellids, diversity indices and proportion of *Harmonia axyridis* at two different sites in the Federal District during the period of August 2008 to January 2010.

Species	Collection sites		Total
	Gama	Taguatinga	
<i>Azya luteipes</i>	0	7	7
<i>Coleomegilla maculata</i>	1	0	1
<i>Cycloneda sanguinea</i>	50	32	82
<i>Eriopis connexa</i>	199	19	218
<i>Harmonia axyridis</i>	1	109	110
<i>Hippodamia convergens</i>	266	27	293
<i>Hyperaspis festiva</i>	5	6	11
<i>Nephaspis</i> spp.	14	0	14
<i>Olla v-nigrum</i>	0	17	17
<i>Psyllobora</i> spp.	3	21	24
<i>Scymnus</i> spp.	65	39	104
Richness (S)	9	9	
Abundance	604	277	
Dominance Index	0.321	0.212	
Shannon Wiener (H)	1.346	1.846	
Simpson	0.678	0.787	

Species diversity was also higher in Taguatinga than in Gama when compared by the Shannon-Wiener index ($t = -8.045$, $P < 0.001$). The profile of diversity based on the Renyi index (Fig. 5) and the Simpson index

supports this as well, as it reflects that the dominance of the community was higher in Gama (Table 1).

DISCUSSION

We recorded only succinae group individuals, in the same way as that observed by Rezende *et al.* (2010) in Minas Gerais, despite the polymorphism of colors displayed by the species in Brazil. From previous records in Brazil (Almeida & Silva, 2002; Rezende *et al.*, 2010), as well as records in the United States and Mexico (Brown & Miller, 1998), we can infer that this staining pattern is predominant in America. This is in contrast with the Palaearctic region and Europe as a whole where dark shapes are also found (Koch, 2003).

The Gama region was the site where the first specimen of *H. axyridis* was collected on mixed stands of tomato and cilantro in August 2008. According to Osawa (2000), populations of *H. axyridis* show a pattern of dispersal among subpopulations in which their establishment is conditioned by availability and accessibility of aphid colonies. In places such as Gama where there was a low occurrence of aphids, *H. axyridis* may have used this type of habitat as a temporary




FIGURE 3: Seasonal variation in the population of *Harmonia axyridis* and other coccinellid (*Cycloneda sanguinea*, *Hippodamia convergens*, *Eriopis connexa*, *Scymnus* spp. *Azya luteipes*, *Hyperaspis festiva*, *Olla v-nigrum* and *Psyllobora* sp.) collected in an area of vegetable intercropping in organic systems from February 2009 to January 2010 in Taguatinga, Federal District.




FIGURE 4: Estimated richness of species or genera of the coccinellid community through rarefaction curves in vegetable crops in two regions of the Federal District.




FIGURE 5: Diversity profile of coccinellid communities collected among organic vegetable crops in two regions of the Federal District.

refuge in search of better quality patches of resources. The collection of only one specimen, with no observation of any other specimen, is associated with a low occurrence of colonies of aphids in the area. This suggests that their arrival has been recent and migrant populations have not yet been established. In the region of Taguatinga, *H. axyridis* was a constant presence during the collections of coccinellids among vegetables grown in intercropping systems during the year 2009/10. This species presented a seasonal variation related to rainfall distribution in the region and constancy of prey throughout the year, suggesting a possible local establishment. This pattern of seasonal variation according to the region's climate was also observed in the south of the country where the species is already established but where it fluctuates throughout the year (Milléo *et al.*, 2008; Martins *et al.*, 2009).

Considering the previous reports of *H. axyridis* in Brazil, this work may be considered thus far to be the northernmost record of the species in the country. Koch *et al.* (2006) used climate models by comparing climate of the Neotropics with climate of the original areas of *H. axyridis* in the Palearctic region. In this model, authors indicated the south of the country as

the most likely location for establishment, and the Central region of Brazil as a place with low probability for this species to occur. In the Central region, temperatures range between 15° and 30°C and air humidity in the dry season may go as low as 15%, which is a limiting factor for the establishment of this population in the area. However, individuals have been collected during the rainy season and especially during the dry period in the Federal District, demonstrating its great capacity to adapt to new climatic conditions, including areas previously considered unsuitable for its establishment.

After the first record of *H. axyridis* in southern Brazil, the density of this species continued to increase and cause negative impacts on native coccinellid species in the region (Milléo *et al.*, 2008; Martins *et al.*, 2009). Considering the stages of biological invasion, colonization, establishment and spread (Koch *et al.*, 2006), we can infer that this species is already established in Brazil and is in the process of spreading to new areas.

In the Federal District, *H. axyridis* is apparently still in the process of establishment due to low abundance or absence in some sampling sites. Analysis of richness and diversity of the coccinellid community allows us to infer that, despite high relative abundance of *H. axyridis* against coccinellid fauna in Taguatinga, there is no apparent negative impact on diversity of the local community. This inference is based on estimates made in the two sampling areas, showing the greatest richness and diversity of coccinellids in the region of Taguatinga. In this area relative abundance of *H. axyridis* was higher, suggesting that there is still a shift of species as suggested by Milléo *et al.* (2008) and Martins *et al.* (2009) in southern Brazil.

However, relatively low abundance of species in some sampling sites does not exclude the risk of *H. axyridis* becoming a threat to native aphid predators in the future. Since the arrival of the species is apparently new to the region, this work highlights the need to constantly monitor this species' impact on the native community of coccinellids and other aphidofagous insects in the Federal District.

RESUMO

A joaninha asiática, Harmonia axyridis (Pallas, 1773), foi primeiramente registrada no Brasil em 2002 no Estado do Paraná, sendo posteriormente registrada nos estados de São Paulo e Minas Gerais. Essa espécie pode colonizar novas áreas e tornar-se dominante na comunidade local, reduzindo a densidade e diversidade de espécies nativas,

principalmente de coccinélidos. O objetivo deste trabalho foi registrar a presença de *H. axyridis* no Distrito Federal, a flutuação populacional e a sua co-ocorrência com outros coccinélidos. As coletas foram realizadas em uma propriedade rural particular em Taguatinga e no campo experimental da Embrapa Hortaliças no Gama, Distrito Federal, entre agosto/2008-janeiro/2010, utilizando coleta direta dos indivíduos sobre as plantas. Foram coletados 881 coccinélidos, sendo que destes 110 pertencem à espécie *H. axyridis* exclusivamente do grupo *succinea*, nas seguintes plantas: tomate, coentro, milho verde, repolho, couve, couve-flor, brócolis, pepino. Também foi observada a sua co-ocorrência com outras espécies de joaninhas: *Cycloneda sanguinea* (Linnaeus, 1763), *Hippodamia convergens* (Guerin-Meneville 1842), *Eriopis connexa* (German, 1824), *Scymnus* sp., *Nephaspis* sp., *Azya luteipes* (Mulsant, 1850), *Hyperaspis festiva* (Mulsant, 1850), *Olla v-nigrum* (Mulsant, 1866), *Psyllobora* sp. e *Coleomegilla maculata* (De Geer, 1775). Até o momento, não foram registradas interações negativas de *H. axyridis* com essas espécies no Distrito Federal. Além disso, esta região foi apontado anteriormente como tendo uma baixa probabilidade de ocorrência para esta espécie, refletindo *H. axyridis* grande plasticidade adaptativa para novos habitats.

PALAVRAS-CHAVE: Controle Biológico; Diversidade; Espécie exótica; Invasão; Predador.

ACKNOWLEDGEMENTS

The authors thank Dra. Lúcia Massutti de Almeida, Camila Fediuk de Castro, and Geovan Henrique Corrêa from the Universidade Federal do Paraná for identifying specimens of coccinellids, as well as FAPDF and CNPq for financial support.

REFERENCES

- ALMEIDA, L.M. DE & SILVA, V.B. 2002. Primeiro registro de *Harmonia axyridis* (Pallas) (Coleoptera, Coccinellidae): um coccinélido originário da região Paleártica. *Revista Brasileira de Zoologia*, 19:941-944.
- BERKOVENS, N.; BONTE, J.; BERKOVENS, D.; DEFORCE, K.; TIRRY, L. & DE CLERCQ, P. 2008. Pollen as an alternative food for *Harmonia axyridis*. *BioControl*, 53:201-210.
- BROWN, M.W. & MILLER, S.S. 1998. Coccinellidae (Coleoptera) in apple orchards of eastern West Virginia and the impact of invasion by *Harmonia axyridis*. *Entomological News*, 109:136-142.
- BROWN, P.M.J.; ADRIAENS, T.; BATHON, H.; CUPPEN, J.; GOLDARAZENA, A.; HÄGG, T.; KENIS, M.; KLAUSNITZER, B.E.M.; KOVÁR, I.; LOOMANS, A.J.M.; MAJERUS, M.E.N.; NEDVED, O.; PEDERSEN, J.; RABITSCH, W.; ROY, H.E.; TERNOIS, V.; ZAKHAROV, I.A. & ROY, D.B. 2008. *Harmonia axyridis* in Europe: spread distribution of a non-native coccinellid. *BioControl*, 53:5-21.
- BURGIO, G.; SANTI, F. & MAINI, S. 2002. On intra-guild predation and cannibalism in *Harmonia axyridis* (Pallas) and *Adalia bipunctata* L. (Coleoptera: Coccinellidae). *Biological Control*, 24:110-116.
- ELLIOTT, N.; KIECKHEFER, R. & KAUFFMAN, W. 1996. Effects of an invading coccinellid on native coccinellids in an agricultural landscape. *Oecologia*, 105:537-544.
- GREZ, A.; ZAVIEZO, T.; GONZÁLEZ, G. & ROTHMANN, S. 2010. *Harmonia axyridis* in Chile: a new threat. *Ciencia y Investigación Agraria*, 37:145-149.
- GONZÁLEZ, G. 2007. *Los Coccinellidae de Perú*. Available at: <www.coccinellidae.cl/paginasWebPeru/Paginas/InicioPeru.php>. Access 17/05/2011.
- HAMMER, O.; HARPER, D.A.T. & RYAN, P.D. 2001. Paleontological statistics software package for education and data analyses. *Paleontologia Electronica*, 4:1-9.
- IABLOKOFF-KHNZORIAN, S.M. 1982. *Les coccinelles Coléoptères-Coccinellidae: Tribu Coccinellini des régions Paléarctique et Orientale*. Société Nouvelle des Éditions Boubée, Paris. 568p.
- KOCH, R.L. 2003. The multicolored Asian lady beetle, *Harmonia axyridis*: A review of its biology, uses in biological control, and non-target impacts. *Journal of Insect Science*, 3:1-16.
- KOCH, R.L.; VENETTE, R.C. & HUTCHISON, W.D. 2006. Invasions by *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae) in the Western hemisphere: implications for South America. *Neotropical Entomology*, 35:421-434.
- MARTINS, C.B.C.; ALMEIDA, L.M.; ZONTA-DE-CARVALHO, R.C.; CASTRO, C.F.; & PEREIRA, R.A. 2009. *Harmonia axyridis*: a threatto Brazilian Coccinellidae? *Revista Brasileira de Entomologia*, 53:663-671.
- MILLÉO, J.; SOUZA, J.M.T. DE; BARBOLA, I.F. & HUSCH, P.E. 2008. *Harmonia axyridis* em árvores frutíferas e impacto sobre outros coccinélidos predadores. *Pesquisa Agropecuária Brasileira*, 43:537-540.
- OSAWA, N. 2000. Population field studies on the aphidophagous ladybird beetle *Harmonia axyridis* (Coleoptera: Coccinellidae): resource tracking and population characteristics. *Population Ecology*, 42:115-127.
- REZENDE, M.Q.; CAMPOS, J.L.A.; COELHO, L.M.B. & SANTANA, D.L.Q. 2010. Coleoptera, Coccinellidae, *Harmonia axyridis* (Pallas, 1773): new record in Minas Gerais, southeastern Brazil. *Checklist*, 6:465-466.
- SAINI, E.D. 2004. Presencia de *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae) en la provincia de Buenos Aires. Aspectos biológicos y morfológicos. *Revista Investigaciones Agropecuarias*, 33:151-160.
- SANTOS, N.R.P.; SANTOS-CIVIDANES, T.M.; CIVIDANES, F.J.; DOS ANJOS, A.C.R. & OLIVEIRA, L.V.L. 2009. Aspectos biológicos de *Harmonia axyridis* alimentada com duas espécies de presas e predação intraguilida com *Eriopis connexa*. *Pesquisa Agropecuária Brasileira*, 44:554-560.
- ZHANG, Z.Q. 1992. The natural enemies of *Aphis gossypii* Glover (Hom., Aphididae) in China. *Journal of Applied Entomology*, 114:251-262.

Recebido em: 23.05.2011
Aceito em: 02.03.2012
Impresso em: 30.03.2012



Publicado com o apoio financeiro do Programa de Apoio às Publicações Científicas Periódicas da USP

EDITORIAL COMMITTEE

Publisher: Museu de Zoologia da Universidade de São Paulo. Avenida Nazaré, 481, Ipiranga, CEP 04263-000, São Paulo, SP, Brasil.

Editor-in-Chief: Carlos José Einicker Lamas, Serviço de Invertebrados, Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42.494, CEP 04218-970, São Paulo, SP, Brasil. E-mail: editormz@usp.br.

Associate Editors: Mário César Cardoso de Pinna (*Museu de Zoologia, Universidade de São Paulo, Brasil*); Luis Fábio Silveira (*Museu de Zoologia, Universidade de São Paulo, Brasil*); Marcos Domingos Siqueira Tavares (*Museu de Zoologia, Universidade de São Paulo, Brasil*); Sérgio Antonio Vanin (*Instituto de Biociências, Universidade de São Paulo, Brasil*); Hussam El Dine Zaher (*Museu de Zoologia, Universidade de São Paulo, Brasil*).

Editorial Board: Aziz Nacib Ab'Saber (*Universidade de São Paulo, Brasil*); Rüdiger Bieler (*Field Museum of Natural History, U.S.A.*); Walter Antonio Pereira Boeger (*Universidade Federal do*

Paraná, Brasil); Carlos Roberto Ferreira Brandão (*Universidade de São Paulo, Brasil*); James M. Carpenter (*American Museum of Natural History, U.S.A.*); Ricardo Macedo Corrêa e Castro (*Universidade de São Paulo, Brasil*); Mario de Vivo (*Universidade de São Paulo, Brasil*); Marcos André Raposo Ferreira (*Museu Nacional, Rio de Janeiro, Brasil*); Darrel R. Frost (*American Museum of Natural History, U.S.A.*); William R. Heyer (*National Museum of Natural History, U.S.A.*); Ralph W. Holzenthal (*University of Minnesota, U.S.A.*); Adriano Brillhante Kury (*Museu Nacional, Rio de Janeiro, Brasil*); Gerardo Lamas (*Museo de Historia Natural "Javier Prado", Lima, Peru*); John G. Maisey (*American Museum of Natural History, U.S.A.*); Náércio Aquino Menezes (*Universidade de São Paulo, Brasil*); Christian de Muizon (*Muséum National d'Histoire Naturelle, Paris, France*); Nelson Papavero (*Universidade de São Paulo, Brasil*); James L. Patton (*University of California, Berkeley, U.S.A.*); Richard O. Prum (*University of Kansas, U.S.A.*); Olivier Rieppel (*Field Museum of Natural History, U.S.A.*); Miguel Trefaut Urbano Rodrigues (*Universidade de São Paulo, Brasil*); Randall T. Schuh (*American Museum of Natural History, U.S.A.*); Ubirajara Ribeiro Martins de Souza (*Universidade de São Paulo, Brasil*); Paulo Emílio Vanzolini (*Universidade de São Paulo, Brasil*); Richard P. Vari (*National Museum of Natural History, U.S.A.*).

INSTRUCTIONS TO AUTHORS - (April 2007)

General Information: *Papéis Avulsos de Zoologia (PAZ)* and *Arquivos de Zoologia (AZ)* cover primarily the fields of Zoology, publishing original contributions in systematics, paleontology, evolutionary biology, ontogeny, faunistic studies, and biogeography. *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* also encourage submission of theoretical and empirical studies that explore principles and methods of systematics.

All contributions must follow the International Code of Zoological Nomenclature. Relevant specimens should be properly curated and deposited in a recognized public or private, non-profit institution. Tissue samples should be referred to their voucher specimens and all nucleotide sequence data (aligned as well as unaligned) should be submitted to GenBank (www.ncbi.nih.gov/Genbank) or EMBL (www.ebi.ac.uk).

Peer Review: All submissions to *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* are subject to review by at least two referees and the Editor-in-Chief. All authors will be notified of submission date. Authors may suggest potential reviewers. Communications regarding acceptance or rejection of manuscripts are made through electronic correspondence with the first or corresponding author only. Once a manuscript is accepted providing changes suggested by the referees, the author is requested to return a revised version incorporating those changes (or a detailed explanation of why reviewer's suggestions were not followed) within fifteen days upon receiving the communication by the editor.

Proofs: Page-proofs with the revised version will be sent to e-mail the first or corresponding author. Page-proofs *must be returned to the editor, preferentially within 48 hours*. Failure to return the proof promptly may be interpreted as approval with no changes and/or may delay publication. Only necessary corrections in proof will be permitted. Once page proof is sent to the author, further alterations and/or significant additions of text are permitted only at the author's expense or in the form of a brief appendix (note added in proof).

Submission of Manuscripts: Manuscripts should be sent to the **SciELO Submission** (<http://submission.scielo.br/index.php/paz/login>), along with a submission letter explaining the importance and originality of the study. Address and e-mail of the corresponding author must be always updated since it will be used to send the 50 reprints in titled by the authors. Figures, tables and graphics **should not** be inserted in the text. Figures and graphics should be sent in separate files with the following formats: ".JPG" and ".TIF" for figures, and ".XLS" and ".CDR" for graphics, with 300 DPI of minimum resolution. Tables should be placed at the end of the manuscript.

Manuscripts are considered on the understanding that they have not been published or will not appear elsewhere in substantially the same or abbreviated form. The criteria for acceptance of articles are: quality and relevance of research, clarity of text, and compliance with the guidelines for manuscript preparation.

Manuscripts should be written preferentially in English, but texts in Portuguese or Spanish will also be considered. Studies with a broad coverage are encouraged to be submitted in English. All manuscripts should include an abstract and key-words in English and a second abstract and key-words in Portuguese or Spanish.

Authors are requested to pay attention to the instructions concerning the preparation of the manuscripts. Close adherence to the guidelines will expedite processing of the manuscript.

Manuscript Form: Manuscripts should not exceed 150 pages of double-spaced, justified text, with size 12 and source Times New Roman (except for symbols). Page format should be A4 (21 by 29.7 cm), with 3 cm of margins. The pages of the manuscript should be numbered consecutively.

The text should be arranged in the following order: **Title Page, Abstracts with Key-Words, Body of Text, Literature Cited, Tables, Appendices, and Figure Captions**. Each of these sections should begin on a new page.

(1) **Title Page:** This should include the **Title, Short Title, Author(s) Name(s) and Institutions**. The title should be concise and, where appropriate, should include mention of families and/or higher taxa. Names of new taxa should not be included in titles.

(2) **Abstract:** All papers should have an abstract in **English** and another in **Portuguese or Spanish**. The abstract is of great importance as it may be reproduced elsewhere. It should be in a form intelligible if published alone and should summarize the main facts, ideas, and conclusions of the article. Telegraphic abstracts are strongly discouraged. Include all new taxonomic names for referencing purposes. Abbreviations should be avoided. It should not include references. Abstracts and key-words should not exceed 350 and 5 words, respectively.

(3) **Body of Text:** The main body of the text should include the following sections: **Introduction, Material and Methods, Results, Discussion, Conclusion, Acknowledgments, and References at end**. Primary headings in the text should be in capital letters, in bold and centered. Secondary headings should be in capital and lower case letters, in bold and centered. Tertiary headings should be in capital and lower case letters, in bold and indented at left. In all the cases the text should begin in the following line.

(4) **Literature Cited:** Citations in the text should be given as: Silva (1998) *or* Silva (1998:14-20) *or* Silva (1998: figs. 1, 2) *or* Silva (1998a, b) *or* Silva & Oliveira (1998) *or* (Silva, 1998) *or* (Rangel, 1890; Silva & Oliveira, 1998a, b; Adams, 2000) *or* (Silva, *pers. com.*) *or* (Silva *et al.*, 1998), the latter when the paper has three or more authors. The reference need not be cited when authors and date are given only as authority for a taxonomic name.

(5) **References:** The literature cited should be arranged strictly alphabetically and given in the following format:

- **Journal Article** - Author(s). Year. Article title. *Journal name*, volume: initial page-final page. Names of journals must be spelled out in full.
- **Books** - Author(s). Year. *Book title*. Publisher, Place.
- **Chapters of Books** - Author(s). Year. Chapter title. *In: Author(s) ou Editor(s), Book title*. Publisher, Place, volume, initial page-final page.
- **Dissertations and Theses** - Author(s). Year. *Dissertation title*. (Ph.D. Dissertation). University, Place.
- **Electronic Publications** - Author(s). Year. *Title*. Available at: <electronic address>. Access in: date.

Tables: All tables must be numbered in the same sequence in which they appear in text. Authors are encouraged to indicate where the tables should be placed in the text. They should be comprehensible without reference to the text. Tables should be formatted with vertical (portrait), not horizontal (landscape), rules. In the text, tables should be referred as Table 1, Tables 2 and 4, Tables 2-6. Use "TABLE" in the table heading.

Illustrations: Figures should be numbered consecutively, in the same sequence that they appear in the text. Each illustration of a composite figure should be identified by capital letters and referred in the text as: Fig. 1A, Fig. 1B, for example. When possible, letters should be placed in the left lower corner of each illustration of a composite figure. Hand-written lettering on illustrations is unacceptable. Figures should be mounted in order to minimize blank areas between each illustration. Black and white or color photographs should be digitized in high resolution (300 DPI at least). Use "Fig(s)," for referring to figures in the text, but "FIGURE(S)" in the figure captions and "fig(s)," when referring to figures in another paper.

Responsibility: Scientific content and opinions expressed in this publication are sole responsibility of the respective authors.
Copyrights: The journals *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* are licensed under a Creative Commons Licence (<http://creativecommons.org>).

For other details of manuscript preparation of format, consult the CBE Style Manual, available from the Council of Science Editors (www.councilscienceeditors.org/publications/style).
Papéis Avulsos de Zoologia and *Arquivos de Zoologia* are publications of the Museu de Zoologia da Universidade de São Paulo (www.mz.usp.br).
Always consult the Instructions to Authors printed in the last issue or in the electronic home pages: www.scielo.br/paz or www.mz.usp.br/publicacoes.